

This quiz assesses your understanding of the articles in this issue. Participation is open to physicians who practise in Canada. Physicians who complete the quiz will receive a statement from the University of Calgary, Office of Continuing Medical Education, indicating their participation, score and the correct answer to each question.

Each quiz may be submitted only once within six months of publication. Correct answers will be

published six months after the quiz appears.

You will receive your score and CME statement immediately if you answer the quiz online at either www.cme.ucalgary.ca or www.stacommunications.com.

A question may have more than one correct answer. Please use a ballpoint pen to mark the correct answers on the quiz reply form. You can complete the quiz online or mark all the correct answers on the quiz reply form.

Select the best answer(s) for each of the following:

1. *What is the half-life of flurazepam?*

- a) 8 to 200 hours
- b) 26 to 200 hours
- c) 30 to 200 hours
- d) 50 to 250 hours
- e) 96 to 250 hours

2. *Which drugs does the author recommend avoiding in the elderly?*

- a) Amitriptyline
- b) Desipramine
- c) Metoclopramide
- d) Piroxicam
- e) Procyclidine

(Drugs and the Elderly, page 41)

3. *Which medications are mentioned in the article as having interactions with celecoxib?*

- a) Fluconazol
- b) Fluvastatine
- c) Lithium
- d) Rifampin
- e) Zafirlukast

(Cox-2 Inhibitors, page 51)

4. *Which route of hormone replacement therapy causes a lower incidence of migraine attacks without aura?*

- a) Oral
- b) Transdermal

(HRT, page 57)

5. *Which of the following are local factors that delay wound healing?*

- a) Arterial insufficiency
- b) Edema
- c) Hyperglycemia
- d) Infection
- e) Venous insufficiency

6. *Chronic wounds should be cleansed with antiseptics.*

- a) True
- b) False

(Chronic Wound Care, page 67)

7. *Which class of the New York Heart Association functional classification is associated with marked limitation of physical activity?*

- a) Class I
- b) Class II
- c) Class III
- d) Class IV
- e) Class V

CME Credit Quiz

8. Following treatment of congestive heart failure with ACE inhibitors, a rise in serum creatinine of 20 % is a reason to discontinue treatment.

- a) True
- b) False

(Cardiac Insufficiency, page 73)

9. Which drugs are serotonin-noradrenalin modulators?

- a) Bupropion
- b) Citalopram
- c) Mirtazapine
- d) Moclobemide
- e) Venlafaxine

(Depression, page 78)

10. Clopidogrel is the recommended initial anti-platelet therapy in cases of peripheral arterial disease?

- a) True
- b) False

11. Which stage of the Fontaine Classification is associated with claudication pain at rest?

- a) Stage I
- b) Stage II
- c) Stage III
- d) Stage IV
- e) Stage V

12. There is no current data to justify using an angiotensin receptor antagonist for preventing systemic atherosclerotic complications in patients with peripheral arterial disease.

- a) True
- b) False

(Peripheral Arterial Disease, page 85)

13. Establishing a no-suicide contract is the first step in managing a potential suicide.

- a) True
- b) False

(Youth at Risk, page 93)

If you answer the quiz online at www.stacommunications.com or www.cme.ucalgary.ca, you will receive your score and CME statement immediately.

If you do not have Internet access, fax the answer sheet to **(403) 270-7285**. If you do not have a fax machine, mail it to:

**CME Quiz, c/o Continuing Medical Education
The University of Calgary
3330 Hospital Drive NW
Calgary, Alberta
T2N 4N1**

**Made possible through an educational grant from
MERCK FROSST CANADA LTD.**