

This quiz assesses your understanding of the articles in this issue. Participation is open to physicians who practise in Canada. Physicians who complete the quiz will receive a statement from the University of Calgary, Office of Continuing Medical Education, indicating their participation, score and the correct answer to each question.

Each quiz may be submitted only once within six months of publication. Correct answers will be published six months after the quiz appears.

You will receive your score and CME statement immediately if you answer the quiz online at either www.cme.ucalgary.ca or www.stacommunications.com.

A question may have more than one correct answer. Please use a ballpoint pen to mark the correct answers on the quiz reply form. You can complete the quiz online or mark all the correct answers on the quiz reply form.

Select the best answer(s) for each of the following:

1. *Which test may detect an Achilles tendon rupture?*

- a) McBurney
- b) MacDermott
- c) McThomas
- d) Smith
- e) Thompson

2. *An Achilles tendon can function with only 25 % of its fibres intact.*

- a) True
- b) False

3. *After the rupture or repair of an Achilles tendon, how long should a patient avoid rebounding activities?*

- a) 4 weeks
- b) 6 weeks
- c) 8 weeks
- d) 10 weeks
- e) 12 weeks

(Ankle Injuries, page 43)

4. *Incontinence is a normal part of aging.*

- a) True
- b) False

5. *What is the lifetime risk of infection with Helicobacter pylori infection in third-world countries?*

- a) 10 %
- b) 30 %
- c) 50 %
- d) 70 %
- e) 90 %

6. *According to Laheij et al, which treatment regimen for Helicobacter pylori has the lowest cure percentage?*

- a) AMTP
- b) BAM
- c) PAC
- d) PCM
- e) RAC

(H. pylori, page 53)

7. *Migraines are more common in men than in women.*

- a) True
- b) False

8. *Cluster headaches are more common in women than in men.*

- a) True
- b) False

CME Credit Quiz

9. What is the recommended treatment for paroxysmal hemicrania?

- a) Caffeine
- b) Codeine
- c) Ibuprofen
- d) Indomethacin
- e) Lithium

10. After subarachnoid hemorrhage, how long does the cerebrospinal fluid (CSF) xanthochromia remain positive?

- a) 1-2 days
- b) 3-5 days
- c) 6-9 days
- d) 10-14 days
- e) 15-21 days

(Headaches, page 83)

11. In the National Institutes of Health (NIH) Stroke Scale, what score is assigned for complete paralysis of one or both sides of the face?

- a) 1
- b) 2
- c) 3
- d) 4
- e) 5

12. In the NIH Stroke Scale assessment of language, what score is assigned for severe aphasia?

- a) 1
- b) 2
- c) 3
- d) 4
- e) 5

(Stroke, page 91)

13. What percentage of cases of von Willebrand's disease is Type I?

- a) 5 %
- b) 15 %
- c) 33 %
- d) 85 %
- e) 95 %

14. Mild von Willebrand's disease is an autosomal recessive disorder.

- a) True
- b) False

(Bleeding Disorders, page 101)

If you answer the quiz online at www.stacommunications.com or www.cme.ucalgary.ca, you will receive your score and CME statement immediately.

If you do not have Internet access, fax the answer sheet to (403) 270-7285. If you do not have a fax machine, mail it to: **CME Quiz, c/o Continuing Medical Education**

**The University of Calgary
3330 Hospital Drive NW
Calgary, Alberta
T2N 4N1**

**Made possible through an educational grant from
MERCK FROSST CANADA LTD.**