

This quiz assesses your understanding of the articles in this issue. Participation is open to physicians who practise in Canada. Physicians who complete the quiz will receive a statement from the University of Calgary, Office of Continuing Medical Education, indicating their participation, score and the correct answer to each question.

Each quiz may be submitted only once within six months of publication. Correct answers will be

published six months after the quiz appears.

You will receive your score and CME statement immediately if you answer the quiz online at either www.cme.ucalgary.ca or www.stacommunications.com.

A question may have more than one correct answer. Please use a ballpoint pen to mark the correct answers on the quiz reply form. You can complete the quiz online or mark all the correct answers on the quiz reply form.

Select the best answer(s) for each of the following:

1. *Less than half the patients with invasive meningococcal disease will have a rash.*

- a) True
- b) False

2. *Which drugs are suggested for the first line treatment of bacterial meningitis?*

- a) Ampicilin
- b) Cefotaxime
- c) Ceftriaxone
- d) Rifampin
- e) Vancomycin

(Meningitis and Encephalitis, page 43)

3. *What percentage of 10-month-old infants will have gastroesophageal reflux?*

- a) 5% to 10%
- b) 20% to 30%
- c) 30% to 40%
- d) 40% to 50%
- e) 50% to 60%

4. *Which symptoms are side-effects of bethanecol therapy?*

- a) Agitation
- b) Cramps
- c) Diarrhea
- d) Insomnia
- e) Urine incontinence

(GI Reflux in Children, page 53)

5. *Before what age should symptoms have been present, when considering a diagnosis of ADD?*

- a) 2
- b) 4
- c) 7
- d) 14
- e) 21

6. *What percentage of children with ADD will have a parent with the disorder?*

- a) 2%
- b) 5%
- c) 10%
- d) 15%
- e) 20%

(ADD in Adults, page 64)

7. *Approximately what percentage of active tuberculosis cases are picked up via X-ray?*

- a) 70%
- b) 75%
- c) 80%
- d) 85%

8. *Culture-and-sensitivity-directed therapy usually includes three active agents for two months, followed by _____.*

- a) Four active agents for two months
- b) Three active agents for four months
- c) Two active agents for four months
- d) Four active agents for three months

(Tuberculosis, page 73)

CME Credit Quiz

9. *What percentage of circulating testosterone is produced in the ovarian stroma?*

- a) 5%
- b) 10%
- c) 25%
- d) 50%
- e) 75%

10. *What per cent of testosterone is bound to sex hormone binding globulin?*

- a) 2%
- b) 32%
- c) 45%
- d) 66%
- e) 98%

(Low Libido in Women, page 83)

11. *What is the common denominator between anorexia nervosa and bulimia?*

- a) Absence of menses
- b) Binge eating
- c) Distorted body image
- d) Fear of obesity
- e) Nausea

12. *Which conditions are organic causes of chronic nausea or vomiting?*

- a) Brain tumour
- b) Cyclic vomiting syndrome
- c) Diabetic gastroparesis
- d) Hyperthyroidism
- e) Intestinal pseudo-obstruction

(GI Dysmotility in Adolescence, page 91)

13. *Which "A" word is not part of the "three As" cited in the palliative care article?*

- a) Attentiveness
- b) Apathy
- c) Accompaniment
- d) Advocacy

14. *Non-oncology patients generally differ in _____ ways that will influence orders and approaches from symptom control.*

- a) Three
- b) Four
- c) Five
- d) Six

(Palliative Care, page 99)

If you answer the quiz online at www.stacommunications.com or www.cme.ucalgary.ca, you will receive your score and CME statement immediately.

If you do not have Internet access, fax the answer sheet to **(403) 270-7285**. If you do not have a fax machine, mail it to:

**CME Quiz, c/o Continuing Medical Education
The University of Calgary
3330 Hospital Drive NW
Calgary, Alberta
T2N 4N1**

**Made possible through an educational grant from
MERCK FROSST CANADA LTD.**